

Comment to the reply to "On the thermodynamics of inhomogeneous perfect fluid mixtures"

Bartolomé Coll¹ and Joan Josep Ferrando²

¹ Systèmes de référence relativistes, SYRTE, Observatoire de Paris-CNRS, 75005 Paris, France.

² Departament d'Astronomia i Astrofísica, Universitat de València, 46100 Burjassot, València, Spain.

E-mail: bartolome.coll@obspm.fr; joan.ferrando@uv.es

Abstract. In spite of a recent reply by Quevedo and Zárate (gr-qc/0403096), their assertion that their thermodynamic scheme for a perfect fluid binary mixture is incompatible with Szekeres and Stephani families of universes, except those of FRW ones, remains wrong.

In the paper "Thermodynamic scheme of inhomogeneous perfect fluid mixtures" [1], Quevedo and Zárate introduce a thermodynamic scheme for a binary mixture of relativistic perfect fluids with non-vanishing entropy production, and analyze its compatibility with the Szekeres and Stephani families of perfect fluid solutions to Einstein Equations.

Their thermodynamic scheme is very clearly presented, the two families of solutions are well known from long time, and in any moment of their argumentation additional restrictions are made due to physical considerations or to supposedly known variables. In this *general* context fixed by them, they conclude that their thermodynamic scheme is incompatible with *all* the space-times of both families of solutions except the Friedmann-Robertson-Walker ones.

This conclusion of incompatibility, repeatedly asserted in their paper [1], is wrong.

In our Comment "On the thermodynamics of inhomogeneous perfect fluid mixtures" [2], we proved the wrongness of their assertion by *three independent* ways:

i) giving the correct analysis of compatibility and proving that, contrarily to their assertion, not only *all* the perfect fluid space-times admit their scheme, but that every one of them admits *many* different such schemes,

ii) showing the errors in two of their crucial arguments that lead them to their false conclusion and

iii) giving simple counterexamples to their assertion.

Surprisingly enough, their Reply [3] to our Comment:

i') only presents vague comments about *imprecise* 'physical meanings' and 'known variables', not referred to in their original paper, which anyway are *strongly insufficient*

to constitute a set of forgotten or implicit conditions in [1] that could be able to transform their wrong assertion into a correct one.

ii') does not consider the two errors in their crucial arguments that lead them to their false conclusion,

iii') does not takes into account the counterexamples to their assertion.

As a consequence of these three points (but only one of them suffices!), *the mentioned assertion, even to the light of their replay [3], remains clearly wrong*, as indeed may be verified by any advanced student in the subject.

Our intention has not been, and is not, polemic. Essentially we try to avoid other searches, particularly those non specialists but interested in the subject, to lose their time (and money ?) studying or assuming such a wrong result.

For this reason and the strong evidence of the errors in paper [1], from now on we shall not comment on any other vague argumentation on this subject.

NOTES

1. In their Reply [3] it is asserted that the conclusion of our Comment [2] is erroneous because based on an erroneous assumption (namely, that the chemical potential and fractional concentration of a mixture of perfect fluids are unknown variables). This assertion is *impertinent* because *hypotheses non fécimus* in our Comment, at least neither more nor less than them in their original paper [1].

2. In their Reply [3] it is asserted that in our Comment [2] we apply inappropriately the 'Pfaffian method' to a 'non Pfaffian' form. This assertion is also *impertinent* because the only result from J. F. Pfaff that we use is his local decomposition theorem for *arbitrary* one-forms, later recovered by G. Frobenius, G. Darboux and E. Cartan among others. Although one may well be unaware of these historical details, our description of the contents of the theorem ('local decomposition'), our explicit expression (5) and the simplicity of the context makes hard to understand that the word 'Pfaff' could so perturb and so confuse the authors of the Reply [3]. It is obvious that, contrarily to their assertion, the (Pfaff) decomposition theorem used in our Comment is not only *completely* appropriate but is *the* appropriate one to be applied.

3. It remains, as it is evident, that the two families of solutions in question cannot admit arbitrarily given binary mixtures. The problem of finding all the (formally) admissible binary mixtures, and that of finding, among them, those which may be physically interesting for every particular application remain open. Nevertheless, and as already stated in our Comment [2] (last paragraph of Section 2), the richness of the choice of Zárate and Quevedo's schemes stated in our Proposition 2 locally guarantees the usual thermodynamic inequalities (such as $T > 0$, or $0 \leq c \leq 1$).

[1] Zárate R D and Quevedo H 2004 *Class. Quantum Grav.* **21** 197 *Preprint* gr-qc/0310087

[2] Coll B and Ferrando J J 2003 *Preprint* gr-qc/0312058

[3] Quevedo H and Zárate R D 2004 *Preprint* gr-qc/0403096